

Fermat's Principle

Last time, talked about light scattering:

total field = incident field + scattered field

Transmission through media:

- scattered field causes phase shift,
looks like wave slows down

But individually, both waves travel at c

1

Today: Start considering what happens at boundary between two materials

Get Law of Reflection, Snell's Law
Generalize to Fermat's Principle

Outline:

- Optical materials
- Scattering and reflection
- Refraction
- Fermat's principle

Next time: finish studying boundaries

2

Optical Materials

Talked about index and absorption:

Said good materials have no resonances in visible

Be a little more specific

What material to use in given application?

References:

- Optics catalogs (Melles Griot, CVI, Oriel)
- Schott Glass catalog

3

Most common optical glass: Schott BK7

SiO_2 with B_2O_5 , Na_2O , CaO + others

Index $n \approx 1.5$


Transmission range 350 – 2000 nm

Few bubbles or defects

Use for windows, lenses in visible, near-infrared

4

Schott BK7 Glass



5

Resonance in UV:
electronic excitations

Resonance in IR:
molecular excitations

Impurities: several small resonances, $\lambda = 1-2 \mu\text{m}$
don't see on graph
important for lasers, optical fibers

Question: Why does absorption increase so much more slowly in IR than in UV?

6

Other useful materials:

Schott SF11 glass: transmits 400 nm – 2 μm
 $n \approx 1.7$

Pyrex: good mirror substrate

Suprasil: transmits 150 nm – 2 μm

MgF₂: transmits 150 nm – 6 μm
used in coatings

CaF₂: transmits 150 nm – 9 μm

Sapphire: transmits 200 nm – 6 μm

ZeSe: transmits 700 nm – 20 μm

Many others

7

Reflection (Hecht 4.3)

Again, two approaches possible:


- Use Maxwell with $\epsilon_0 \rightarrow \epsilon$
- Think about scattered fields

Today: take scattering approach
try to understand physics

Next time: use Maxwell, get complete answers

8

Start with light in piece of glass:




Why no reflection from say $z = 0$?

Scattered waves from nearby atoms cancel out

9

Introduce gap at $z = 0$:



Missing atoms: scattered waves don't cancel

Remove material, reflected wave appears!

10

Expect wave from both surfaces

Fields should be equal and opposite:

would cancel if surfaces brought together

Funny point:

Does reflection comes from atoms at surface?

No:

Reflection = net wave from all atoms $z > 0$


(Which atoms were missing ones cancelling?)

Question: If reflected light comes from all the atoms, shouldn't the reflected field change if you introduce a second surface downstream? How can that be?

11

Usually think about atoms on surface for simplicity

In general, incident light at an angle:




Expect reflected wave where all scattered fields have same phase

12

Say plane waves from distant source α ,
 detect at distance point β

Consider scattered field from atoms A and B



13

Need fields from A and B to have same phase

Say distance from α to A = L

distance from B to β = L'

Incident field at A = $E_0 e^{i[kL - \omega t]}$

Incident field at B = $E_0 e^{i[k(L + d_B) - \omega t]}$

Scattered field from A at β

$$A e^{ikL} e^{i[k(L' + d_A) - \omega t]}$$


Scattered field from B at β :

$$A e^{ik(L + d_B)} e^{i[kL' - \omega t]}$$

14

Make phases equal: $L + L' + d_A = L + L' + d_B$

or $d_A = d_B$



angle of incidence θ_i

angle of reflection θ_r


Need $\boxed{\theta_i = \theta_r}$: Law of Reflection

15

Typically just draw k-vectors

= "rays"

and don't think about all this scattering stuff



Just like balls bouncing off a wall

But good to know what's going on underneath

16

Sometimes, need 3D version of reflection law

Define $\hat{\mathbf{u}}$ = normal to surface

Then $\hat{\mathbf{k}}_{\text{refl}}$ in plane of $\hat{\mathbf{k}}_{\text{inc}}$ and $\hat{\mathbf{u}}$

Get


$$\hat{\mathbf{k}}_{\text{refl}} = \hat{\mathbf{k}}_{\text{inc}} - 2(\hat{\mathbf{k}}_{\text{inc}} \cdot \hat{\mathbf{u}})\hat{\mathbf{u}}$$

Question: Should $\hat{\mathbf{u}}$ be normal pointing out of or into the surface?

17

Refraction (Hecht 4.4)

Think about transmitted wave



Now have $\mathbf{E}_{\text{tot}} = \mathbf{E}_{\text{inc}} + \mathbf{E}_{\text{scat}}$

(For reflection, \mathbf{E}_{scat} and \mathbf{E}_{inc} more distinct)

18

But \mathbf{E}_{scat} hard to calculate now:

- Scattered field is strong, gets rescattered
- Want field inside medium, close to charges

Be clever instead:

Incident medium: $n = n_1$

Total incident wave = $\mathbf{E}_1 e^{i(n_1 \mathbf{k}_1 \cdot \mathbf{r} - \omega t)}$

Take $\mathbf{k} = k_{1x} \hat{\mathbf{x}} + k_{1z} \hat{\mathbf{z}}$

Transmitted medium: $n = n_2$

Total transmitted wave = $\mathbf{E}_2 e^{i(n_2 \mathbf{k}_2 \cdot \mathbf{r} - \omega t)}$

Then $\mathbf{k} = k_{2x} \hat{\mathbf{x}} + k_{2z} \hat{\mathbf{z}}$

19

At boundary $z = 0$:

$$\mathbf{E}_{\text{inc}} = \mathbf{E}_1 e^{i(n_1 k_{1x} x - \omega t)} \quad \mathbf{E}_{\text{trans}} = \mathbf{E}_2 e^{i(n_2 k_{2x} x - \omega t)}$$


Don't need \mathbf{E} continuous across boundary,
do need phase difference independent of x

- Since boundary is uniform, phase relationship between waves can't depend on x
- Same reason ω doesn't change, phase difference can't depend on time

So need $n_1 k_{1x} = n_2 k_{2x}$

20

Geometry:



$$k_{1x} = k_1 \sin \theta_1$$

$$k_{2x} = k_2 \sin \theta_2$$

21

Here k_1 and k_2 are vacuum k 's: $k_1 = k_2 = \omega/c$

So if $n_1 k_{1x} = n_2 k_{2x}$

$$\text{then } \frac{n_1 \omega}{c} \sin \theta_1 = \frac{n_2 \omega}{c} \sin \theta_2$$

or

$$\boxed{n_1 \sin \theta_1 = n_2 \sin \theta_2} \quad \text{Snell's Law}$$

Follows from:

- $\mathbf{k} \rightarrow n\mathbf{k}$ in medium
- symmetry of surface

22

Generalize to 3D:

Have \mathbf{k}_{inc} , $\mathbf{k}_{\text{trans}}$ and surface normal $\hat{\mathbf{u}}$
in same plane

Write

$$n_1 \hat{\mathbf{k}}_1 \times \hat{\mathbf{u}} = n_2 \hat{\mathbf{k}}_2 \times \hat{\mathbf{u}}$$

or

$$\mathbf{k}_1 \times \hat{\mathbf{u}} = \mathbf{k}_2 \times \hat{\mathbf{u}}$$

23

Fermat's Principle (Hecht 4.5)

Can generalize previous results

Think about reflection again


Before asked:

- Where does detector need to be to see reflected light?

Now ask:

- Given source and detector positions, which points on surface contribute to detected light?

24




Which atoms on surface are radiating waves that interfere constructively at β ?

Hard to work out geometrically, since phase of \mathbf{E}_{inc} is complicated

(Even worse if surface is curved!)

25

Consider point P on surface



Incident field at $P = E_0 e^{i(k s - \omega t)}$

Field from P reaching β : $E'_0 e^{i[k(s+s') - \omega t]}$

Write $s + s' = S \equiv \text{Optical Path Length}$

26

Net phase at β : $\phi_P = k\mathcal{S}$

Get constructive interference from points near P
if fields from nearby points have same phase

Or: if \mathcal{S} constant near P

If P labelled by coordinate x , want

$$\frac{d\mathcal{S}}{dx} = 0$$

27

Work this out

Say $\mathbf{r}_\alpha = (x_1, z_1)$ and $\mathbf{r}_\beta = (x_2, z_2)$ and $\mathbf{r}_P = (x, 0)$
(surface at $z = 0$, leave out y for now)

$$\begin{aligned}\text{Then } \mathcal{S} &= s + s' \\ &= \sqrt{(x - x_1)^2 + z_1^2} + \sqrt{(x - x_2)^2 + z_2^2}\end{aligned}$$

$$\begin{aligned}\text{and } \frac{d\mathcal{S}}{dx} &= \frac{x - x_1}{\sqrt{(x - x_1)^2 + z_1^2}} + \frac{x - x_2}{\sqrt{(x - x_2)^2 + z_2^2}} \\ &= 0\end{aligned}$$

28

Solve for x :
$$\frac{(x - x_1)^2}{(x - x_1)^2 + z_1^2} = \frac{(x - x_2)^2}{(x - x_2)^2 + z_2^2}$$

Invert:
$$1 + \frac{z_1^2}{(x - x_1)^2} = 1 + \frac{z_2^2}{(x - x_2)^2}$$


So
$$\frac{z_1}{x - x_1} = \pm \frac{z_2}{x - x_2}$$

From original equation, need “-” root

Solve
$$x = \frac{x_1 z_2 + x_2 z_1}{z_1 + z_2}$$

29

Relate to geometry:



Had

$$\frac{x - x_1}{\sqrt{(x - x_1)^2 + z_1^2}} = \frac{x_2 - x}{\sqrt{(x - x_2)^2 + z_2^2}}$$

Means $\sin \theta_i = \sin \theta_r$

Again, $\theta_i = \theta_r$, law of reflection

30

Idea: light takes path such that \mathcal{S} is stationary
 \equiv constant for small variations in path

“Path” identifies atoms whose scattered fields add constructively

Called *Fermat's Principle*

Very powerful method

31


Apply to refraction

Phase from α to β

$$= n_1 k s + n_2 k s'$$

Here define $\mathcal{S} = n_1 s + n_2 s'$

Again set $d\mathcal{S}/dx = 0$



Question: Why do we want $d\mathcal{S}/dx = 0$ here?


32

$$\mathcal{S}(x) = n_1 \sqrt{(x - x_1)^2 + z_1^2} + n_2 \sqrt{(x - x_2)^2 + z_2^2}$$

$$\frac{d\mathcal{S}}{dx} = \frac{n_1(x - x_1)}{s} + \frac{n_2(x - x_2)}{s'} = 0$$

Again, gives

$$n_1 \sin \theta_1 = n_2 \sin \theta_2$$



33

For arbitrary path, define

$$\mathcal{S} = \int n(\mathbf{r}) ds \quad (\text{integrated along path})$$

Usually path = sum of straight line segments

then $\int \rightarrow \sum$

Fermat's Principle:

Light takes path such that \mathcal{S} is *stationary*

Small variations in path, \mathcal{S} doesn't change
(could be min, max, or constant)

Question: What does Fermat's principle say about light travelling through free space?

34

To use:

If \mathcal{S} is function of parameters $\{x_i\}$, want

$$\frac{\partial \mathcal{S}}{\partial x_i} = 0 \quad \text{for all } i$$

For physics students:

Can allow arbitrary path variations

write $\delta \mathcal{S} = 0$,

get differential equation for path

Just like mechanics

35

Note, haven't really proven Fermat's Principle:

- Works for reflection
makes sense from scattering picture
- Works for refraction
scattering picture unclear
- Works in free space
no scattering at all!

Also, ambiguous what “path” means for a wave

Revisit when we derive Huygens' Principle

36

Summary

For now, understand how light reflected, refracted
can understand lenses, mirrors

- Law of reflection: $\theta_i = \theta_r$
- Snell's law: $n_1 \sin \theta_1 = n_2 \sin \theta_2$

- Fermat's principle:

$$\mathcal{S} = \int n \, ds$$

light travels path with \mathcal{S} constant