

A Mini Time Projection Chamber for the PEN Experiment

L. Peter Alonzi III

University of Virginia

April APS Meeting
Washington D.C., 14 February 2010

Outline

PEN Experiment: Motivation and Context

Mini Time Projection Chamber (mTPC)

Global Context $\pi \rightarrow e\nu$

THEORY: $BR = \frac{\Gamma(\pi \rightarrow e\nu(\gamma))}{\Gamma(\pi \rightarrow \mu\nu(\gamma))} =$

$$\left\{ \begin{array}{l} (1.2352 \pm 0.0005) \times 10^{-4} \quad \text{Marciano and Sirlin, [PRL 71 (1993) 3629]} \\ (1.2354 \pm 0.0002) \times 10^{-4} \quad \text{Finkemeier, [Phys. Lett. B 387 (1996) 391]} \\ (1.2352 \pm 0.0001) \times 10^{-4} \quad \text{Cirigliano and Rosell, [PRL 99, 231801 (2007)]} \end{array} \right.$$

EXPERIMENT [PDG]: $BR = (1.230 \pm 0.004) \times 10^{-4}$; $\frac{\delta BR}{BR} \approx 3.3 \times 10^{-3}$

PEN GOAL: $\frac{\delta BR}{BR} < 5 \times 10^{-4}$

Global Context $\pi \rightarrow e\nu$

THEORY: $BR = \frac{\Gamma(\pi \rightarrow e\nu(\gamma))}{\Gamma(\pi \rightarrow \mu\nu(\gamma))} =$

$$\left\{ \begin{array}{ll} (1.2352 \pm 0.0005) \times 10^{-4} & \text{Marciano and Sirlin, [PRL 71 (1993) 3629]} \\ (1.2354 \pm 0.0002) \times 10^{-4} & \text{Finkemeier, [Phys. Lett. B 387 (1996) 391]} \\ (1.2352 \pm 0.0001) \times 10^{-4} & \text{Cirigliano and Rosell, [PRL 99, 231801 (2007)]} \end{array} \right.$$


EXPERIMENT [PDG]: $BR = (1.230 \pm 0.004) \times 10^{-4}$; $\frac{\delta BR}{BR} \approx 3.3 \times 10^{-3}$

PEN GOAL: $\frac{\delta BR}{BR} < 5 \times 10^{-4}$

PEN Experimental Program


Precision measurement of hadronic charged current decay

- $\pi^+ \rightarrow e^+ \nu_e$
 - $e - \mu$ universality
 - **V-A** coupling deviations (pseudoscalar)
 - m_{h^+} , PS/V leptoquarks, etc.
- $\pi^+ \rightarrow e^+ \nu_e \gamma$
 - **V-A** coupling deviations (tensor)
 - F_A/F_V , χ PT test
- $\mu^+ \rightarrow e^+ \nu_e \bar{\nu}_\mu \gamma$
 - \mathcal{L}_{weak} departures from V-A



PEN Experimental Setup


- π E1 beamline at PSI
- Stopped π^+ beam
- 240-module CsI calorimeter
- Active target
- Central tracking
- Some Systematics
 - Decay in flight events
 - CsI gainmatching
 - Detector Alignment







PEN Experimental Setup

- π E1 beamline at PSI
- Stopped π^+ beam
- 240-module CsI calorimeter
- Active target
- Central tracking


- Some Systematics
 - Decay in flight events
 - CsI gainmatching
 - Detector Alignment



Decay In Flight Kinematics





mTPC Technical Specifications





- Proportional Region: 40x6x40 mm
- Drift Region: 40x40x50 mm
- Drift Gas: 90% He and 10% C₂H₆
- 4000 V across drift region
- Grid: 50 μ m wires with 1 mm spacing
- Nichrome Anode Wires
 - 40 mm length
 - 20 μ m diameter
 - 10 mm spacing
 - 235 Ω resistance
- CAEN VME digitizer V1720

Fabricated by our collaborators from Dubna, Russia




Waveform Digitization




- x: charge division
- y: drift time
- z: wire location


mTPC Coordinate Calibration



- MWPC coordinates well known
- Calibrate mTPC with MWPC



mTPC coordinate Resolution




$$\delta_i = i_2 - i_1 + \frac{(i_0 - i_3)}{3}$$

$$\sigma_i = \frac{RMS_{\delta_i}}{\sqrt{1^2 + 1^2 + (1/3)^2 + (1/3)^2}}$$

$$\Rightarrow \sigma_x < 1.3 \text{ mm (charge division)}$$

$$\Rightarrow \sigma_y < 0.35 \text{ mm (drift time)}$$

mTPC coordinate Resolution



$$\delta_i = i_2 - i_1 + \frac{(i_0 - i_3)}{3}$$


$$\sigma_i = \frac{RMS_{\delta_i}}{\sqrt{1^2 + 1^2 + (1/3)^2 + (1/3)^2}}$$

$\Rightarrow \sigma_x < 1.3 \text{ mm}$ (charge division)

$\Rightarrow \sigma_y < 0.35 \text{ mm}$ (drift time)

Results from 2009 Data Run

Decay Vertex Prediction



- X Position
 - charge division
 - $\sigma_x < 1.3$ mm

- Y position
 - drift time
 - $\sigma_y < 0.35$ mm

Experiment R-05-01 (PEN) collaboration members:

L. P. Alonzi^a, V. A. Baranov^b, W. Bertl^c, M. Bychkov^a, Yu.M. Bystritsky^b, E. Frlež^a, V.A. Kalinnikov^b, N.V. Khomutov^b, A.S. Korenchenko^b, S.M. Korenchenko^b, M. Korolija^d, T. Kozlowski^e, N.P. Kravchuk^b, N.A. Kuchinsky^b, M.C. Lehman^a, D. Mekterović^d, D. Mzhavia^{b,f}, A. Palladino^{a,c}, D. Počanić^{a*}, P. Robmann^g, A.M. Rozhdestvensky^b, S.N. Shkarovskiy^b, U. Straumann^g, I. Supek^d, P. Truöl^g, Z. Tsamalaidze^f, A. van der Schaaf^{g*}, E.P. Velicheva^b, and V.P. Volnykh^b

^aDept of Physics, Univ. of *Virginia*, Charlottesville, VA 22904-4714, USA

^bJoint Institute for Nuclear Research, RU-141980 *Dubna*, Russia

^cPaul Scherrer Institut, CH-5232 Villigen *PSI*, Switzerland

^dInstitut "Rudjer Bošković", HR-10000 *Zagreb*, Croatia

^eInst. Problemów Jądrowych im. Andrzeja Sołtana PL-05-400 *Swierk*, Poland

^fIHEP, Tbilisi State University, GUS-380086 *Tbilisi*, Georgia

^gPhysik Institut der Universität *Zürich*, CH-8057 Zürich, Switzerland

Web page: <http://pen.phys.virginia.edu>